

Dark energy and the future fate of the Universe

Yungui Gong

*College of Electronic Engineering, Chongqing University of Posts and Telecommunications,
Chongqing 400065, China*

gongyg@cqupt.edu.cn

Yuan-Zhong Zhang

*CCAST (World Laboratory), P.O. Box 8730, Beijing 100080 and
Institute of Theoretical Physics, Chinese Academy of Sciences, P.O. Box 2735, Beijing 100080,
China*

We consider the possibility of observing the onset of the late time inflation of our patch of the Universe. The Hubble size criterion and the event horizon criterion are applied to several dark energy models to discuss the problem of future inflation of the Universe. We find that the acceleration has not lasted long enough to confirm the onset of inflation by present observations for the dark energy model with constant equation of state, the holographic dark energy model and the generalized Chaplygin gas (GCG) model. For the flat Λ CDM model with $\Omega_{m0} = 0.3$, we find that if we use the Hubble size criterion, we need to wait until the a_v which is the scale factor at the time when the onset of inflation is observed reaches 3.59 times of the scale factor a_T when the Universe started acceleration, and we need to wait until $a_v = 2.3a_T$ to see the onset of inflation if we use the event horizon criterion. For the flat holographic dark energy model with $d = 1$, we find that $a_v = 3.46a_T$ with the Hubble horizon and $a_v = 2.34a_T$ with the event horizon, respectively. For the flat GCG model with the best supernova fitting parameter $\alpha = 1.2$, we find that $a_v = 5.50a_T$ with the Hubble horizon and $a_v = 2.08a_T$ with the event horizon, respectively.

Keywords: dark energy; future fate.

PACS Nos.: 04.20.Gz, 98.80.Cq

1. Introduction

There are increasing evidences that the Universe is expanding with acceleration and the transition from decelerated expansion to accelerating expansion happened in the recent past. The transition redshift $z_T > 0.2$, we use the subscript T to denote the transition throughout this paper. These results suggest that there exists dark energy (DE) with negative pressure in the Universe, and the DE was subdominant in the past and dominates the Universe now. The presence of DE has a lot of interesting physical effects. For example, there exists an event horizon if the acceleration is eternal. The event horizon sets a causal limit that the observers can ever access. The existence of eternal acceleration also prevents us from ever measuring inflationary perturbations which originated before the ones currently observable^{1,2}.

The DE physics is still a challenging topic. The current supernova Ia (SN Ia) data is unable to distinguish different DE models and different DE parameterizations ^{3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18}. Starkman, Trodden and Vachaspati (STV) addressed the problem of inflation in our patch of the Universe with the help of the concept of the minimal anti-trapped surface (MAS) ¹⁹. They argued that if we can confirm the acceleration up to a redshift z_c and observe the contraction of our MAS, then we are certain that our universe is inflating. If we see the contraction of our MAS, then we observe the onset of inflation. The immediate conclusion is that our universe is undergoing inflation because the cosmic acceleration is confirmed up to the redshift 1.755 by the SN 1997ff. STV found that the period of acceleration has not lasted long enough for observations to confirm the onset of inflation for the Λ CDM model.

The work of STV is based on the earlier work of Vachaspati and Trodden, who proved that in a homogeneous and isotropic universe, the necessary and sufficient condition for observing the contraction of the MAS is that the Universe is vacuum dominated in a region of radius greater than the Hubble size H^{-1} ²⁰. The comoving contraction of our MAS is the essence of inflation. Thus only if a region of size greater than H^{-1} remains vacuum dominated long enough for the MAS to begin collapsing then we are certain that the Universe is undergoing inflation ¹⁹. Because the Hubble size is increasing with time in general, so the later the transition time (the smaller the redshift z_T) is, the longer inflation needs to last (the larger a_v/a_T is, where a_v is the scale factor at the time when the onset of acceleration is first seen.). Avelino, de Carvalho and Martins then replaced the Hubble size by the event horizon with some additional assumptions ²¹. If the event horizon criterion is used, then the smaller z_T is, the smaller a_v/a_T we need. Huterer, Starkman and Trodden also analyzed general DE models and found that current observations are unable to confirm the onset of inflation ²². In this paper, we discuss the holographic DE model ^{23,24,25,26,27,28} and the generalized Chaplygin gas (GCG) model ^{29,30,31,32}.

2. The Hubble Size Criterion

Vachaspati and Trodden proved that inflationary models based on the classical Einstein equations, the weak energy conditions, and trivial topology, must assume homogeneity on super-Hubble scales. Based on this result, STV introduced the concept of the MAS to discuss the observability of the onset of inflation. The MAS is a sphere, centered on the observer, on which the velocity of comoving objects is the speed of light c ¹⁹. For light emitted directly toward the observer inside the MAS, the photons get closer to the observer with time, while all photons emitted by sources outside the MAS get farther away. For a homogeneous and isotropic universe, the physical radius of the MAS at time t is the Hubble size $1/H(t)$. It was argued that the beginning of the comoving contraction of the MAS can be identified with the onset of inflation. Note that the condition of the onset of inflation is $\ddot{a} = 0$ which is equivalent to $d(aH)^{-1}/dt$, where $a(t)$ is the scale factor.

If a light was emitted at time t_e from a source located at a comoving distance r and then is received at time t_v by the observer located at the origin, the physical distance between the source and the observer at t_e is

$$d(t_e, t_v) = a(t_e) \int_{t_e}^{t_v} \frac{dt}{a(t)}. \quad (1)$$

In this paper, we consider a flat universe. We can see the contraction of the MAS at time t_v if $d(t_e, t_v) = 1/H(t_e)$ ¹⁹, here t_v (a_v) is the time (scale factor) when the turnaround of the MAS comes into view. Therefore, the onset of the late time inflation can be seen at time t_v if $d(t_T, t_v) = 1/H(t_T)$, where t_T is the transition time when the Universe experienced from the deceleration phase to the acceleration phase. To see the consequences of $d(t_e, t_v) = 1/H(t_e)$ clearly, we use a simple DE model $p = w\rho$ with constant w satisfying $w \geq -1$ as an example.

The matter energy density is $\rho_m = \rho_{mr}(a_r/a)^3$, where the subscript r means that the variable takes a value at an arbitrary reference time t_r . The DE density is $\rho_x = \rho_{xr}(a_r/a)^{3(1+w)}$. So we have

$$H^2 = H_r^2 \left[\Omega_{mr} \left(\frac{a_r}{a} \right)^3 + \Omega_{xr} \left(\frac{a_r}{a} \right)^{3(1+w)} \right], \quad (2)$$

where $\Omega = 8\pi G\rho/(3H^2)$. The transition time t_T is determined from

$$\Omega_{mT} + (1 + 3w)\Omega_{xT} = 0, \quad (3)$$

or

$$1 + z_T = \frac{a_0}{a_T} = \left(-\frac{\Omega_{m0}}{\Omega_{x0}(1 + 3w)} \right)^{1/3w}, \quad (4)$$

where $\Omega_{m0} = 1 - \Omega_{x0}$ and the subscript 0 means that the variable takes its present value. For the Λ CDM model, $w = -1$ and $\Omega_{m0} = 0.3$, we get $z_T = 0.67$. Using Eq. (3), the condition $d(t_T, t_v) = 1/H(t_T)$ becomes

$$\int_{a_T/a_v}^1 \frac{dx}{\sqrt{x^3(x^{3w} - 1 - 3w)}} = \frac{1}{\sqrt{-3w}}. \quad (5)$$

To be able to observe the onset of inflation at present, we require $a_v < a_0$. For the Λ CDM model, the solution to the above equation (5) is $a_v/a_T = 3.59 > 1 + z_T = 1.67$, so $a_v > a_0$ and $\Omega_{\Lambda v} = 0.96$. In addition, we require the confirmation of cosmic acceleration up to a redshift z_c , where z_c determined from the condition $d(t_c, t_0) = 1/H(t_c)$ satisfies the following equation

$$\int_0^{z_c} \frac{dz}{\sqrt{\Omega_{m0}(1+z)^3 + \Omega_{x0}(1+z)^{3(1+w)}}} = \frac{1 + z_c}{\sqrt{\Omega_{m0}(1+z_c)^3 + \Omega_{x0}(1+z_c)^{3(1+w)}}}. \quad (6)$$

Note that z_c is the minimum redshift that the cosmic acceleration needs to be observed by current observations. For the Λ CDM model with $\Omega_{m0} = 0.3$, we get $z_c = 1.61$. Since the current SN Ia observations extend to redshift 1.755, so the cosmic acceleration is confirmed. But we still need to wait until Ω_{Λ} reaches a value




Fig. 1. The dependence of a_T/a_0 and a_T/a_v on w and Ω_{x0} for the DE model with constant equation of state parameter w . The line labeled “Hubble Horizon” shows the dependence of a_T/a_v on w by using the Hubble scale criterion. The line labeled “Event Horizon” shows the dependence of a_T/a_v on w by using the event horizon criterion.

of 0.96 to observe the onset of inflation. The solutions to Eqs. (4) and (5) for other choices of Ω_{x0} and w are shown in Fig 1. From Fig. 1, we see that we need to wait until $\Omega_x \sim 0.9$ for $w > -0.67$ to be able to observe the onset of inflation. For bigger w , we need smaller Ω_x . However, current observations strongly constrain $w \lesssim -0.8$. In other words, we are unable to confirm the onset of inflation now with current observations.

In Ref. 33, Lightman and Press introduced the concept of constant redshift surfaces to discuss the causal communication of comoving particles. In the definition of the constant redshift surfaces, the lower integral t_e changes with the upper integral t_v in equation (1) by fixing $a(t_v)/a(t_e) = z_1$ to be a constant. The constant redshift surface or z_1 -surface increases with time before inflation. After inflation, the z_1 -surface will eventually decrease with time. For small redshift $z_1 < z_T$, the z_1 -surface is decreasing. For large redshift, the z_1 -surface is increasing. So there exists a turnaround redshift z_1 so that the z_1 -surface reaches its maximum at present. For the Λ CDM model, we find that the turnaround redshift $z_1 = 2.09$ if we take $\Omega_{m0} = 0.3$. Although the decrease of the z_1 -surface for small z_1 is a characteristic feature of an accelerated universe, it does not mean that the Universe is inflating

and the space-time will evolve into the de-Sitter phase. Only if z_1 is big enough so that the z_1 -surface crosses the event horizon, then we can say that the Universe is inflating. We will discuss this in the next section.

2.1. Holographic DE model

Cohen, Kaplan and Nelson proposed that for any state with energy E in the Hilbert space, the corresponding Schwarzschild radius $R_S \sim E$ is less than than the infrared (IR) cutoff L ²³. Therefore, the maximum entropy is $S_{BH}^{3/4}$. Under this assumption, a relationship between the ultraviolet cutoff $\rho_x^{1/4}$ and the IR cutoff is derived, i.e., $8\pi GL^3 \rho_x/3 \sim L$. Hsu found that the model based on the Hubble scale as the IR cutoff would not give an accelerating universe²⁴. Li then showed that a plausible dark energy is possible by choosing the future event horizon as the IR cutoff²⁵. So the holographic DE density is^{25,26}

$$\rho_x = \frac{3d^2}{8\pi G R_{eh}^2}, \quad (7)$$

where $R_{eh}(t) = d(t, \infty)$ is the event horizon. The equation of state of the holographic DE is

$$w_x = -\frac{1}{3} \left(1 + \frac{2\sqrt{\Omega_x}}{d} \right).$$

Because of some physical constraints on d , we take $d = 1$ for simplicity²⁷. Note that the weak energy condition is satisfied as long as $d^2 \geq \Omega_x$. By using the Friedmann equations, we get

$$\frac{d\Omega_x}{d \ln a} = \Omega_x(1 - \Omega_x)(1 + 2\sqrt{\Omega_x}). \quad (8)$$

The solution to Eq. (8) is

$$\ln \Omega_x - \frac{1}{3} \ln(1 - \sqrt{\Omega_x}) + \ln(1 + \sqrt{\Omega_x}) - \frac{8}{3} \ln(1 + 2\sqrt{\Omega_x}) = \ln \left(\frac{a}{a_r} \right) + y_r, \quad (9)$$

where y_r is determined from the above equation by using Ω_{xr} . From the definition of the holographic DE density (7) and the Friedmann equations, we get

$$R_{eh}(t) = \frac{1}{H_r \sqrt{1 - \Omega_{xr}}} \left(\frac{a(t)}{a_r} \right)^{3/2} \left(\frac{1 - \Omega_x(t)}{\Omega_x(t)} \right)^{1/2}. \quad (10)$$

The transition time t_T is determined from

$$\Omega_{xT} + 2\Omega_{xT} \sqrt{\Omega_{xT}} = 1. \quad (11)$$

So $\Omega_{xT} = 0.432$ and $y_T = -2.215$. Substitute these values to Eq. (9), set $a_r = a_0$ and $\Omega_x = \Omega_{xT}$ and use the best fitting value $\Omega_{xr} = \Omega_{x0} = 0.75$, we get $z_T = 0.72$. Since $d(t_T, t_v) = R_{eh}(t_T) - a_T R_{eh}(t_v)/a_v$, the condition $d(t_T, t_v) = 1/H(t_T)$ becomes

$$\frac{a_v(1 - \Omega_{xv})}{a_T \Omega_{xv}} = \frac{(1 - \sqrt{\Omega_{xT}})^2(1 - \Omega_{xT})}{\Omega_{xT}} = 0.154. \quad (12)$$

Combining Eqs. (9) and (12), we get

$$\ln \Omega_{xv} - \frac{1}{3} \ln(1 - \sqrt{\Omega_{xv}}) + \ln(1 + \sqrt{\Omega_{xv}}) - \frac{8}{3} \ln(1 + 2\sqrt{\Omega_{xv}}) = \ln \left(\frac{0.154\Omega_{xv}}{1 - \Omega_{xv}} \right) + y_T. \quad (13)$$

Combining Eqs. (12) and (13), we get $a_v/a_T = 3.46 > 1 + z_T = 1.72$, so $a_v > a_0$. The redshift z_c satisfies the equation

$$\frac{(1 - \sqrt{\Omega_{xc}})^2(1 - \Omega_{xc})}{(1 + z_c)\Omega_{xc}} = \frac{1 - \Omega_{x0}}{\Omega_{x0}}. \quad (14)$$

Combining Eqs. (9) and (14) with $\Omega_{x0} = 0.75$, we get $z_c = 1.64 < 1.755$. Therefore although we see our MAS today, we are unable to observe the onset of inflation for the holographic DE model because the cosmic acceleration has not lasted long enough.

2.2. GCG Model

For the GCG model, we have $p_g = -A/\rho_g^\alpha$. By using the energy conservation equation, we get

$$\rho_g = \rho_{gr} \left[-w_{gr} + (1 + w_{gr}) \left(\frac{a_r}{a} \right)^{3(1+\alpha)} \right]^{1/(1+\alpha)}, \quad (15)$$

where the equation of state parameter $w_g = p_g/\rho_g$. Because $\rho_g \sim (a_0/a)^3$ when $a \ll a_0$ and $\rho_g \sim \text{constant}$ when $a \gg a_0$, the GCG model can be thought as a unified model of DE and dark matter. Therefore we assume that there is no matter present for simplicity and require $w_g \geq -1$ so that the weak energy condition is satisfied. As discussed in Ref. 32, some reasonable physical constraints also require $\alpha \geq 0$. The Friedmann equation is

$$H^2 = H_r^2 \left[-w_{gr} + (1 + w_{gr}) \left(\frac{a_r}{a} \right)^{3(1+\alpha)} \right]^{1/(1+\alpha)}. \quad (16)$$

At the transition time t_T , we have $w_{gT} = -1/3$. So the transition redshift satisfies

$$1 + z_T = \left(-\frac{2w_{g0}}{1 + w_{g0}} \right)^{1/3(1+\alpha)}. \quad (17)$$

By using the best supernova fitting values $w_{g0} = -0.83$ and $\alpha = 1.20$, we get $z_T = 0.412$. The condition $d(t_T, t_v) = 1/H(t_T)$ gives

$$\int_{a_T/a_v}^1 \left[\frac{1}{3} + \frac{2}{3}x^{3(1+\alpha)} \right]^{-1/2(1+\alpha)} dx = 1. \quad (18)$$

If we take $\alpha = 1.2$, we get $a_v/a_T = 5.50$ and $a_v > a_0$. The condition $d(t_c, t_0) = 1/H(t_c)$ gives

$$\int_0^{z_c} \left[-w_{g0} + (1 + w_{g0})(1 + z)^{3(1+\alpha)} \right]^{-1/2(1+\alpha)} dz = \frac{1 + z_c}{\left[-w_{g0} + (1 + w_{g0})(1 + z_c)^{3(1+\alpha)} \right]^{1/2(1+\alpha)}}. \quad (19)$$




Fig. 2. The dependence of a_T/a_0 and a_T/a_v on w_{g0} and α for the GCG model. The line labeled “Hubble Horizon” shows the dependence of a_T/a_v on α by using the Hubble scale criterion. The line labeled “Event Horizon” shows the dependence of a_T/a_v on α by using the event horizon criterion.

By using the best supernova fitting values $w_{g0} = -0.83$ and $\alpha = 1.20$, we get $z_c = 1.424 < 1.755$. Again the present observations of cosmic acceleration extend to a redshift $z_c = 1.424$. Currently we are still unable to confirm the onset of inflation for the GCG model. For some other values of w_{g0} and α , the numerical solutions to Eqs. (17) and (18) are shown in Fig. 2. From Fig. 2, we see that it is possible to confirm the onset of inflation for the GCG model only if $\alpha < 0$ which is outside the physical parameter space.

3. The Event Horizon Criterion

In general, the Hubble size increases with time. Therefore, it will take longer time to observe the onset of inflation if the transition happened at later time. In order to avoid this situation, Avelino, de Carvalho and Martins replaced the Hubble scale criterion discussed in the previous section by requiring that the comoving distance equals to the comoving event horizon at the time of reception. Of course, some additional assumptions on the content of the local universe and field dynamics are

needed. The event horizon criterion is

$$r = \int_{t_e}^{t_v} \frac{dt}{a(t)} = \int_{t_v}^{\infty} \frac{dt}{a(t)}. \quad (20)$$

By using the notation of the constant redshift surface in ³³, the above condition gives us the redshift z_1 when the z_1 -surface crosses the event horizon. To illustrate the effect of this condition, we take the simple DE model with constant equation of state as an example. Applying the condition (20) to the onset of inflation, Eq. (5) is replaced by

$$\int_{a_T/a_v}^1 \frac{dx}{\sqrt{x^3(x^{3w} - 1 - 3w)}} = \int_0^{a_T/a_v} \frac{dx}{\sqrt{x^3(x^{3w} - 1 - 3w)}}, \quad (21)$$

and Eq. (6) is replaced by

$$\int_0^{z_c} \frac{dz}{\sqrt{\Omega_{m0}(1+z)^3 + \Omega_{x0}(1+z)^{3(1+w)}}} = \int_{-1}^0 \frac{dz}{\sqrt{\Omega_{m0}(1+z)^3 + \Omega_{x0}(1+z)^{3(1+w)}}}. \quad (22)$$

For the Λ CDM model, the solution to Eq. (21) is $a_v/a_T = 2.30 > 1 + z_T = 1.67$ or $\Omega_{\Lambda v} = 0.86$, and the solution to Eq. (22) is $z_c = 1.81$ if we take $\Omega_{m0} = 0.3$. So we are unable to confirm the onset of inflation now with current observations. The numerical solutions to Eq. (21) for other values of w are shown in Fig 1. From Fig. 1, we see that we need to wait until $\Omega_x \sim 0.9$ for $w < -0.85$ to be able to observe the onset of inflation. The smaller w is, the sooner we observe the onset of inflation. It is possible that we observe the onset of inflation with $\Omega_{\Lambda 0} = 0.7$ if $w < -1$. However, the weak energy condition is violated if $w < -1$, and the criterions discussed in this paper do not apply. This situation needs to be investigated more carefully and are out of the scope of our discussion.

3.1. Holographic DE model

Applying the event horizon criterion (20) to the holographic DE model discussed in the previous section, we replace Eq. (12) by

$$\frac{a_v(1 - \Omega_{xv})}{a_T \Omega_{xv}} = \frac{1 - \Omega_{xT}}{4\Omega_{xT}} = 0.329, \quad (23)$$

and Eq. (14) by

$$\frac{1 - \Omega_{xc}}{4(1 + z_c)\Omega_{xc}} = \frac{1 - \Omega_{x0}}{\Omega_{x0}}. \quad (24)$$

Combining Eqs. (9) and (23), we get $a_v/a_T = 2.34 > 1 + z_T = 1.72$, so $a_v > a_0$. Combining Eqs. (9) and (24) with $\Omega_{x0} = 0.75$, we get $z_c = 1.84$. Therefore we are unable to observe the onset of inflation now for the holographic DE model.

3.2. GCG Model

Applying the event horizon criterion (20) to the GCG model discussed in the previous section, we replace Eq. (18) by

$$\int_{a_T/a_v}^1 \left[\frac{1}{3} + \frac{2}{3}x^{3(1+\alpha)} \right]^{-1/2(1+\alpha)} dx = \int_0^{a_T/a_v} \left[\frac{1}{3} + \frac{2}{3}x^{3(1+\alpha)} \right]^{-1/2(1+\alpha)} dx, \quad (25)$$

and Eq. (19) by

$$\int_0^{z_c} \frac{dz}{[-w_{g0} + (1 + w_{g0})(1 + z)^{3(1+\alpha)}]^{1/2(1+\alpha)}} = \int_{-1}^0 \frac{dz}{[-w_{g0} + (1 + w_{g0})(1 + z)^{3(1+\alpha)}]^{1/2(1+\alpha)}}. \quad (26)$$

If we take $\alpha = 1.2$, we get $a_v/a_T = 2.08$, so $a_v > a_0$. By using $w_{g0} = -0.83$ and $\alpha = 1.20$, the solution to Eq. (26) is $z_c = 1.64$. Because $a_v > a_0$, currently we are unable to confirm the onset of inflation for the GCG model. For different values of α , the numerical solutions to Eq. (25) are shown in Fig. 2. From Fig. 2, we see that it is possible to confirm the onset of inflation for the GCG model when $\alpha < 0.5$ and $w_{g0} \sim -0.95$. The smaller α is, the smaller w_{g0} is required to observe the onset of inflation.

4. Discussion

For the Hubble size criterion, our results are: (1) Constant $w \geq -1$ model. It is possible to observe the onset of inflation when $\Omega_x \sim 0.9$ and $w > -0.67$. (2) The holographic DE model with $d = 1$. We find that $z_T = 0.72$ and $z_c = 1.64$ if we use the best supernova fitting result $\Omega_{x0} = 0.75$. We also get $a_v/a_T = 3.46$ and $a_v > a_0$. (3) The GCG model. We find that $z_T = 0.412$ and $z_c = 1.424$ if we use the best supernova fitting results $w_{g0} = -0.83$ and $\alpha = 1.20$. By using $\alpha = 1.20$, we get $a_v/a_T = 5.50$ and $a_v > a_0$. It is possible to observe the onset of inflation when $\alpha < 0$ and $w_g \sim -0.75$.

For the event horizon criterion, our results are: (1) Constant $w \geq -1$ model. It is possible to observe the onset of inflation when $\Omega_x \sim 0.9$ and $w < -0.85$. (2) The holographic DE model with $d = 1$. We find that $z_c = 1.84$ if we use the best supernova fitting result $\Omega_{x0} = 0.75$. We also get $a_v/a_T = 2.34$ and $a_v > a_0$. (3) The GCG model. We find that $z_c = 1.64$ if we use the best supernova fitting results $w_{g0} = -0.83$ and $\alpha = 1.20$. By using $\alpha = 1.20$, we get $a_v/a_T = 2.08$ and $a_v > a_0$. It is possible to observe the onset of inflation when $\alpha < 0.5$ and $w_g \sim -0.95$.

In general, the event horizon criterion gives bigger value of z_c than the Hubble size criterion does. The reason is that today we may be able to observe a larger portion of the Universe than that we can ever access. The event horizon criterion is not applicable for very low values of the dark energy density. However, the later the cosmic acceleration started, the sooner we are able to observe the onset of inflation by using the event horizon criterion.

For all the three models discussed in this paper, we have $z_c > 1$ and $a_v > a_0$. Therefore it is impossible to confirm the onset of inflation by current observations for all three models. However, this conclusion cannot be extended to phantom models because they violate the weak energy condition. Therefore, the conclusion cannot apply to the holographic DE model with phantom behavior³⁴. However, it can be applied to the interacting holographic DE model discussed in Refs. 35 and 36 because $w_{tot} \geq -1$. The conclusion is neither applicable to the GCG model with $w_g < -1$ discussed in Ref. 37. When we parameterized DE equation of state, we find that the supernova data might not be able to distinguish those parameterizations that have almost the same past behaviors and different future behaviors¹⁵. For general dark energy model, it is impossible that cosmic acceleration started at a redshift $z_T > 1$, so it is impossible to observe the onset of inflation up to a region of Hubble size. If the event horizon criterion is used, then it is possible to observe the onset of inflation and confirm the inflation of our universe for some general dynamic dark energy models with low transition redshift. In this paper, we find that the acceleration has not lasted long enough for observations to confirm that we are undergoing inflation. Therefore the future fate of the Universe is still unknown from current observations. We need to wait some time to be confident that we are undergoing inflation. On the other hand, if the cosmic acceleration never ends, less information about the early inflationary perturbations will be observed in the future. So we are living in a peculiar era in the history of the Universe.

For the GCG model, the marginal allowed parameter spaces $\alpha \lesssim 0.5$ and $w_{g0} \gtrsim -0.75$ will make it possible to confirm that our universe is inflating by using the event horizon criterion.

Acknowledgments

Y. Gong is supported by NNSFC under grant No. 10447008 and 10605042, SRF for ROCS, State Education Ministry, CQMEC under grant No. KJ060502. Y.Z. Zhang's work was in part supported by NNSFC under Grant No. 90403032 and also by National Basic Research Program of China under Grant No. 2003CB716300.

References

1. D.A. Lowe and D. Marolf, Phys. Rev. D **70**, 026001 (2004).
2. N. Kaloper, M. Kleban and L. Sorbo, Phys. Lett. B **600**, 7 (2004).
3. A.G. Riess *et al.*, Astrophys. J. **607**, 665 (2004).
4. U. Alam, V. Sahni, T.D. Saini and A.A. Starobinsky, Mon. Not. Roy. Astron. Soc. **354**, 275 (2004).
5. U. Alam, V. Sahni and A.A. Starobinsky, J. Cosmol. Astropart. Phys., JCAP **0406** (2004) 008.
6. R.A. Daly and S.G. Djorgovski, Astrophys. J. **597**, 9 (2003).
7. R.A. Daly and S.G. Djorgovski, Astrophys. J. **612**, 652 (2004).
8. S. Hannestad and E. Mortsell, J. Cosmol. Astropart. Phys., JCAP **0409** (2004) 001.

9. H.K. Jassal, J.S. Bagla and T. Padmanabhan, Mon. Not. Roy. Astron. Soc. **356**, L11 (2005).
10. T.R. Choudhury and T. Padmanabhan, Astron. Astrophys., **429**, 807 (2005).
11. H.K. Jassal, J.S. Bagla and T. Padmanabhan, Phys. Rev. D **72**, 103503 (2005).
12. R. Lazkoz, S. Nesseris and L. Perivolaropoulos, J. Cosmol. Astropart. Phys., JCAP 0511 (2005) 010.
13. Y.G. Gong, Int. J. Mod. Phys. D **14**, 599 (2005).
14. Y.G. Gong, Class. Quantum Grav. **22**, 2121 (2005).
15. Y.G. Gong and Y.Z. Zhang, Phys. Rev. D **72**, 043518 (2005).
16. Y.G. Gong and A. Wang, Phys. Rev. D **73**, 083506 (2006).
17. Y.G. Gong and A. Wang, Phys. Rev. D **75**, 043520 (2007).
18. R. Colistete Jr. and J.C. Fabris, Class. Quantum Grav. **22**, 2813 (2005).
19. G. Starkman, M. Trodden and T. Vachaspati, Phys. Rev. Lett. **83**, 1510 (1999).
20. T. Vachaspati and M. Trodden, Phys. Rev. D **61**, 023502 (2000).
21. P.P. Avelino, J.P.M. de Carvalho and C.J.A.P. Martins, Phys. Lett. B **501**, 257 (2001).
22. D. Huterer, G.D. Starkman and M. Trodden, Phys. Rev. D **66**, 043511 (2002).
23. A. Cohen, D. Kaplan and A. Nelson, Phys. Rev. Lett. **82**, 4971 (1999).
24. S.D.H. Hsu, Phys. Lett. B **594**, 13 (2004).
25. M. Li, Phys. Lett. B **603**, 1 (2004).
26. Q.G. Huang and Y.G. Gong, J. Cosmol. Astropart. Phys., JCAP **0408** (2004) 006.
27. Y.G. Gong, B. Wang and Y.Z. Zhang, Phys. Rev. D **72**, 043510 (2005).
28. Y.G. Gong and Y.Z. Zhang, Class. Quantum Grav. **22**, 4895 (2005).
29. A. Kamenshchik, U. Moschella and V. Pasquier, Phys. Lett. B **511**, 265 (2001).
30. M. Bento, O. Bertolami and A.A. Sen, Phys. Rev. D **66**, 043507 (2002).
31. M.C. Bento, O. Bertolami and A.A. Sen, Phys. Rev. D **70**, 083519 (2004).
32. Y.G. Gong, J. Cosmol. Astropart. Phys., JCAP **0503** (2005) 007.
33. A.P. Lightman and W.H. Press, Astrophys. J. **337**, 598 (1989).
34. S. Nojiri and S.D. Odintsov, Gen. Rel. Grav. **38**, 1285 (2006).
35. D. Pavón and W. Zimdahl, Phys. Lett. B **628**, 206 (2005).
36. B. Wang, Y.G. Gong and E. Abdalla, Phys. Lett. B **624**, 141 (2005).
37. A.A. Sen and R.J. Scherrer, Phys. Rev. D **72**, 063511 (2005).